



FDA Grants Approval for Use of RISPERDAL® CONSTA® as Both a Monotherapy and Adjunctive Therapy in the Maintenance Treatment of Bipolar I Disorder

RISPERDAL® CONSTA® is the First and Only Long-Acting Antipsychotic Therapy Available for Bipolar I Disorder

Titusville, N.J., May 18, 2009 - Janssen®, Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc. today announced the U.S. Food and Drug Administration (FDA) has approved the Supplemental New Drug Applications (sNDAs) for the use of RISPERDAL® CONSTA® (risperidone) Long-Acting Treatment as both monotherapy and adjunctive therapy to lithium or valproate in the maintenance treatment of Bipolar I Disorder.

Bipolar Disorder is a brain disorder that causes unusual shifts in a person's mood, energy and ability to function. It is often characterized by debilitating mood swings from extreme highs (mania) to extreme lows (depression). Type I Bipolar Disorder is characterized based on the occurrence of at least one manic episode, with or without the occurrence of a major depressive episode, and affects approximately 1 percent of the American adult population in any given year.

"Long-acting therapies are moving to the forefront of treatment for mental illness, and the approval of risperidone long-acting treatment for Bipolar Disorder is exciting because it offers physicians assurance that the medication is being taken as prescribed," said Caleb Adler, M.D., principal investigator and associate professor of Clinical Psychiatry at the University of Cincinnati. "Further the bi-weekly administration schedule encourages regular contact between patients and their treatment team."

The approval is based on two prospective, randomized, double-blind, placebo-controlled studies for the long-term treatment of Bipolar I Disorder. The first demonstrated that RISPERDAL® CONSTA®, when used as a monotherapy, was significantly better than placebo at delaying the time to relapse of any mood episode. The second study demonstrated that, for patients already taking lithium or valproate, the addition of RISPERDAL® CONSTA® significantly delayed the time to relapse compared to current treatments plus placebo.

"We are very pleased with this FDA approval for RISPERDAL® CONSTA®," said Hussein Manji, M.D., Global Therapeutic Area Head, Neuroscience, Johnson & Johnson Pharmaceutical Research and Development. "We are committed to creating new, long-acting therapies that offer safe and effective products for treating patients with mental illnesses. This approval provides physicians and patients with a new treatment option that offers a convenient and effective choice to delay relapse."

RISPERDAL® CONSTA® was approved in 2003 as an atypical antipsychotic agent indicated for the treatment of schizophrenia and is now the first and only long-acting injectable antipsychotic therapy available for the treatment of schizophrenia and Bipolar I Disorder.

Visit <http://www.risperdalconsta.com> for full prescribing information.

About RISPERDAL® CONSTA®

Risperidone long-acting treatment (RLAT) is a long-acting injectable form of risperidone that was developed utilizing Alkermes' proprietary Medisorb® drug-delivery technology. Using this technology, risperidone is encapsulated in microspheres made of a biodegradable polymer, which are suspended in a water-based solution and administered to patients by intramuscular injection once every two weeks. RLAT is manufactured by Alkermes, Inc. and marketed by Janssen, Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc. in the U.S. and Janssen-Cilag outside of the U.S.

About J&JPRD

Johnson & Johnson Pharmaceutical Research & Development, L.L.C. (J&JPRD) is a member of the Johnson & Johnson family of companies, the world's most broadly-based producer of health care products. J&JPRD is headquartered in Raritan, N.J., and has facilities throughout Europe, the United States and Asia. J&JPRD is leveraging drug discovery and drug development in a variety of therapeutic areas, including CNS, Internal Medicine and Oncology, to address unmet medical needs worldwide. More information can be found at <http://www.jnjpharmarnd.com>.

About Janssen

Janssen, Division of Ortho-McNeil-Janssen Pharmaceuticals, Inc., is based in Titusville, N.J. and is the only large

pharmaceutical company in the U.S. dedicated solely to mental health. It currently markets prescription medications for the treatment of schizophrenia, bipolar mania and the treatment of symptoms associated with autistic disorder. Ortho-McNeil-Janssen Pharmaceuticals, Inc. is a member of the Johnson & Johnson family of companies. For more information about Janssen[®], visit <http://www.janssen.com/>.

IMPORTANT SAFETY INFORMATION FOR CONSUMERS ABOUT RISPERDAL[®] CONSTA[®]

Elderly Patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death compared to placebo. RISPERDAL[®] CONSTA[®] (risperidone) is not approved for the treatment of patients with dementia-related psychosis.

Neuroleptic Malignant Syndrome (NMS) is a rare and potentially fatal side effect reported with RISPERDAL[®] CONSTA[®] and similar medicines. Call your doctor immediately if the person being treated develops symptoms such as high fever; stiff muscles; shaking; confusion; sweating; changes in pulse, heart rate, or blood pressure; or muscle pain and weakness. Treatment should be stopped if the person being treated has NMS.

Tardive Dyskinesia (TD) is a serious, sometimes permanent side effect reported with RISPERDAL[®] CONSTA[®] and similar medications. TD includes uncontrollable movements of the face, tongue, and other parts of the body. The risk of developing TD and the chance that it will become permanent is thought to increase with the length of therapy and the overall dose taken by the patient. This condition can develop after a brief period of therapy at low doses, although this is much less common. There is no known treatment for TD, but it may go away partially or completely if therapy is stopped.

High blood sugar and diabetes have been reported with RISPERDAL[®] CONSTA[®] and similar medications. If the person being treated has diabetes or risk factors such as being overweight or a family history of diabetes, blood sugar testing should be performed at the beginning and throughout treatment with RISPERDAL[®] CONSTA[®]. Complications of diabetes can be serious and even life threatening. If signs of high blood sugar or diabetes develop, such as being thirsty all the time, going to the bathroom a lot, or feeling weak or hungry, contact your doctor.

RISPERDAL[®] CONSTA[®] and similar medications can raise the blood levels of a hormone known as prolactin, causing a condition known as hyperprolactinemia. Blood levels of prolactin remain elevated with continued use. Some side effects seen with these medications include the absence of a menstrual period; breasts producing milk; the development of breasts by males; and the inability to achieve an erection. The connection between prolactin levels and side effects is unknown.

Some people taking RISPERDAL[®] CONSTA[®] may feel faint or lightheaded when they stand up or sit up too quickly. By standing up or sitting up slowly and following your healthcare professional's dosing instructions, this side effect can be reduced or it may go away over time.

RISPERDAL[®] CONSTA[®] may affect your alertness or driving ability; therefore, do not drive or operate machinery before talking to your healthcare professional.

RISPERDAL[®] CONSTA[®] should be used cautiously in people with a seizure disorder, who have had seizures in the past, or who have conditions that increase their risk for seizures.

Extrapyramidal Symptoms (EPS) are usually persistent movement disorders or muscle disturbances, such as restlessness, tremors, and muscle stiffness. If you observe any of these symptoms, talk to your healthcare professional.

Inform your healthcare professional if you become pregnant or intend to become pregnant during therapy with RISPERDAL[®] CONSTA[®]. Caution should be exercised when RISPERDAL[®] CONSTA[®] is administered to a nursing woman.

RISPERDAL[®] CONSTA[®] may make you more sensitive to heat. You may have trouble cooling off, or be more likely to become dehydrated, so take care when exercising or when doing things that make you warm.

Some medications interact with RISPERDAL[®] CONSTA[®]. Please inform your healthcare professional of any medications or supplements that you are taking. Avoid alcohol while on RISPERDAL[®] CONSTA[®].

RISPERDAL[®] CONSTA[®].

In a study of people taking RISPERDAL[®] CONSTA[®], the most common side effects in the treatment of schizophrenia were

headache, tremors, dizziness, restlessness, tiredness, constipation, indigestion, sleepiness, weight gain, pain in the limbs, and dry mouth.

If you have any questions about RISPERDAL[®] CONSTA[®] or your therapy, talk with your doctor.

IMPORTANT SAFETY INFORMATION FOR PROFESSIONALS ABOUT RISPERDAL[®] CONSTA[®]

WARNING: Increased Mortality in Elderly Patients with Dementia-Related Psychosis Elderly patients with dementia-related psychosis treated with antipsychotic drugs are at an increased risk of death. Analyses of 17 placebo-controlled trials (modal duration of 10 weeks), largely in patients taking atypical antipsychotic drugs, revealed a risk of death in the drug-treated patients of between 1.6 to 1.7 times the risk of death in placebo-treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. Observational studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality. The extent to which the findings of increased mortality in observational studies may be attributed to the antipsychotic drug as opposed to some characteristic(s) of the patients is not clear. RISPERDAL[®] CONSTA[®] (risperidone) is not approved for the treatment of patients with dementia-related psychosis.

Cerebrovascular Adverse Events (CAEs): CAEs, including fatalities, have been reported in elderly patients with dementia-related psychosis taking oral risperidone in clinical trials. The incidence of CAEs with risperidone was significantly higher than with placebo. RISPERDAL[®] CONSTA[®] is not approved for the treatment of patients with dementia-related psychosis.

Neuroleptic Malignant Syndrome (NMS): NMS, a potentially fatal symptom complex, has been reported with the use of antipsychotic medications, including RISPERDAL[®] CONSTA[®]. Clinical manifestations include muscle rigidity, fever, altered mental status and evidence of autonomic instability (see full Prescribing Information). Management should include immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy, intensive symptomatic treatment and medical monitoring, and treatment of any concomitant serious medical problems.

Tardive Dyskinesia (TD): TD is a syndrome of potentially irreversible, involuntary, dyskinetic movements that may develop in patients treated with antipsychotic medications. The risk of developing TD and the likelihood that dyskinetic movements will become irreversible are believed to increase with duration of treatment and total cumulative dose. Elderly patients appeared to be at increased risk for TD. Prescribing should be consistent with the need to minimize the risk of TD. The syndrome may remit, partially or completely, if antipsychotic treatment is withdrawn.

Hyperglycemia and Diabetes: Hyperglycemia, some cases extreme and associated with ketoacidosis, hyperosmolar coma or death has been reported in patients treated with atypical antipsychotics (APS), including RISPERDAL[®] CONSTA[®]. Patients starting treatment with APS who have or are at risk for diabetes should undergo fasting blood glucose testing at the beginning of and during treatment. Patients who develop symptoms of hyperglycemia should also undergo fasting blood glucose testing.

Hyperprolactinemia: As with other drugs that antagonize dopamine D₂ receptors, RISPERDAL[®] CONSTA[®] elevates prolactin levels and the elevation persists during chronic administration. Risperidone is associated with higher levels of prolactin elevation than other antipsychotic agents.

Orthostatic Hypotension: RISPERDAL[®] CONSTA[®] may induce orthostatic hypotension associated with dizziness, tachycardia, and in some patients, syncope, especially during the initial dose-titration period. Monitoring should be considered in patients for whom this may be of concern. RISPERDAL[®] CONSTA[®] should be used with caution in patients with known cardiovascular disease, and conditions that would predispose patients to hypotension.

Leukopenia, Neutropenia and Agranulocytosis have been reported with antipsychotics, including risperidone. Patients with a pre-existing low white blood cell count (WBC) or a history of leukopenia/neutropenia should have frequent complete blood cell counts during the first few months of therapy. At the first sign of a decline in WBC and in the absence of other causative factors, discontinuation of Risperdal Consta should be considered.

Potential for Cognitive and Motor Impairment: RISPERDAL[®] CONSTA[®] has the potential to impair judgment, thinking, or motor skills. Patients should be cautioned about operating hazardous machinery, including motor vehicles, until they are reasonably certain that RISPERDAL[®] CONSTA[®] does not affect them adversely.

Seizures: RISPERDAL[®] CONSTA[®] should be used cautiously in patients with a history of seizures or with conditions that potentially lower seizure threshold.

Dysphagia: Esophageal dysmotility and aspiration can occur. Use cautiously in patients at risk for aspiration pneumonia.

Priapism has been reported. Severe priapism may require surgical intervention.

Thrombotic Thrombocytopenic Purpura (TTP) has been reported.

Administration: Care should be taken to avoid inadvertent injection into a blood vessel.

Suicide: The possibility of suicide attempt is inherent in psychotic illnesses. Close supervision of high-risk patients should accompany drug therapy.

Increased sensitivity in patients with Parkinson's disease or those with dementia with Lewy bodies has been reported. Manifestations and features are consistent with NMS.

Use Risperdal Consta with caution in patients with conditions and medical conditions that could affect metabolism or hemodynamic responses. (e.g. Recent Myocardial infarction or unstable cardiac disease)

Extrapyramidal Symptoms (EPS): The overall incidence of EPS-related adverse events in patients treated with 25 mg and 50 mg of RISPERDAL[®] CONSTA[®] and placebo, respectively, were akathisia* (4%, 11%, 6%), Parkinsonism? (8%, 15%, 9%) and tremor (0%, 3%, 0%).

* Akathisia and restlessness

? Extrapyramidal disorder, musculoskeletal stiffness, muscle rigidity, and bradykinesia

Weight Gain: In a 12-week trial, the percentage of patients experiencing weight gain (>7% of baseline body weight) was 6% placebo versus 9% RISPERDAL[®] CONSTA[®].

Maintenance Treatment: Patients should be periodically reassessed to determine the need for continued treatment.

Commonly Observed Adverse Reactions for RISPERDAL[®] CONSTA[®]: The most common adverse reactions in clinical trials in patients with schizophrenia (75%) were headache, Parkinsonism, dizziness, akathisia, fatigue, constipation, dyspepsia, sedation, weight increase, pain in extremities, and dry mouth.

The most common adverse reactions in clinical trials in patients with bipolar disorder trials were weight increased (5% in monotherapy trial) and tremor and parkinsonism (> 10% in adjunctive therapy trial).

(This press release contains "forward-looking statements" as defined in the Private Securities Litigation Reform Act of 1995. These statements are based on current expectations of future events. If underlying assumptions prove inaccurate or unknown risks or uncertainties materialize, actual results could vary materially from Janssen and/or Johnson & Johnson's expectations and projections. Risks and uncertainties include general industry conditions and competition; economic conditions, such as interest rate and currency exchange rate fluctuations; technological advances and patents attained by competitors; challenges inherent in new product development, including obtaining regulatory approvals; domestic and foreign health care reforms and governmental laws and regulations; and trends toward health care cost containment. A further list and description of these risks, uncertainties and other factors can be found in Exhibit 99 of Johnson & Johnson's Annual Report on Form 10-K for the fiscal year ended December 28, 2008. Copies of this Form 10-K, as well as subsequent filings, are available online at www.sec.gov, www.jnj.com or on request from Johnson & Johnson. Neither Janssen nor Johnson & Johnson undertake to update any forward-looking statements as a result of new information or future events or developments.)

Contacts:

Media

Srikant Ramaswami:
Office: (908) 927-7978
Cell: (609) 647-8195
Email: Sramaswa@its.jnj.com

Investors

Louise Mehrotra: (732) 524-6491

Johnson & Johnson

Lesley Fishman: (732) 524-3922

Johnson & Johnson